



**UNIVERSITI PUTRA MALAYSIA**

***AGROBACTERIUM-MEDIATED TRANSFORMATION OF OIL PALM  
(*ELAEIS GUINEENSIS* JACQ.) SUSPENSION CULTURE***

**SITI HABSAH ROOWI**

**FSMB 2003 19**

***AGROBACTERIUM*-MEDIATED TRANSFORMATION OF OIL PALM  
(*ELAEIS GUINEENSIS* JACQ.) SUSPENSION CULTURE**

**By**

**SITI HABSAH ROOWI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in  
Fulfilment of the Requirement for the Degree of Master of Science**

**June 2003**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

***AGROBACTERIUM-MEDIATED TRANSFORMATION OF OIL PALM  
(ELAEIS GUINEENSIS JACQ.) SUSPENSION CULTURE***

**By**

**SITI HABSAH ROOWI**

**June 2003**

**Chairman : Associate Prof Dr. K. Harikrishna**

**Faculty : Food Science and Biotechnology**

The life cycle of flowering plants in general can be divided into two growth phases: vegetative and reproductive. The reproductive phase can be subdivided into the development of the inflorescence meristem and floral meristems. Control of flowering and the regulation of plant architecture has been thoroughly investigated in a number of well-studied dicot plants such as *Arabidopsis*, *Antirrhinum*, tomato and tobacco. However, in monocot plants, molecular information related to plant reproduction is still limited. In

*Arabidopsis*, the *Terminal Flowering 1* (*TFL1*) gene, *LEAFY*, and the target genes of *CONSTANS* (*CO*) including the *FLOWERING LOCUS T* (*FT*) and *SUPPRESSOR OF OVEREXPRESSION OF CO 1* (*SOC1*) genes, have a major role in promoting flowering and thus controlling flowering time.

To investigate the regulation of meristem identity as well as the control of floral transition in oil palm, we transferred genes *pCAMBIA/TFL1*, *pCAMBIA/JIT60*, *pCAMBIA/LFY*, *pGA/LFY* and *pCAMBIA/SOC1* into oil palm embryogenic callus. This present study focuses on the optimization of *Agrobacterium*-mediated transformation and the analysis of transgenic plants. The objective of this study was also to clone the putative oil palm (*OPSOC1* and *OPLFY*) genes into a plasmid binary vector system, so as to transform these genes into oil palm embryogenic callus and to determine their effect on expression driven by a cauliflower mosaic virus (CaMV) 35S promoter in oil palm.

The success of gene delivery using *Agrobacterium* into the plant genome is often based on several factors including temperature used during co-cultivation, the binary vector and promoters used, and the plant genotype itself. The most important factors contributing to the success of T-DNA transfer is the type of plant material used. We used embryogenic suspension cells as starting material for the transformation of oil palm because of the large numbers of totipotent cells found in these cultures.

The results of the present study show that, embryogenic suspension cells are suitable for *Agrobacterium*-mediated transformation of oil palm. Stable *in vitro* transgenic oil palm plants have been successfully produced through *Agrobacterium*-mediated DNA

delivery within 7-8 months after co-cultivation. A total of 101 transgenic plants, each derived from single embryogenic clusters were obtained from 5563 *Agrobacterium* infected embryogenic clusters from four oil palm lines. The overall transformation frequency was about 1.82%, as it is an average frequency across four parameters (preculture of the embryogenic callus, the effect of temperature during co-cultivation, the effect of binary vectors and the effects of genotype on transformation rate), it is likely that a higher transformation frequency could be obtained routinely under a more optimal set of conditions. Analysis using PCR and Southern Blotting confirmed that the T-DNA was successfully integrated into the oil palm genome.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

***AGROBACTERIUM-MEDIATED TRANSFORMATION OF OIL PALM (*ELAEIS GUINEENSIS* JACQ.) SUSPENSION CULTURE***

**Oleh**

**SITI HABSAH ROOWI**

**June 2003**

**Pengerusi : Profesor Madya Dr. K. Harikrishna**

**Fakulti : Sains Makanan dan Bioteknologi**

Kitaran hidup tumbuhan berbunga secara amnya boleh dibahagikan kepada 2 peringkat pertumbuhan : vegetatif dan reproduktif. Fasa reproduktif pula boleh dibahagikan kepada pembentukan meristem infloresen dan meristem bunga. Pengawalan pembungaan dan rupabentuk tumbuhan telah dikaji secara mendalam pada beberapa tumbuhan dikot seperti *Arabidopsis*, *Antirrhinum*, tomato dan tembakau. Walaubagaimanapun, pada tumbuhan monocot, maklumat molekular dalam penghasilan tumbuhan masih terhad. Di dalam tumbuhan *Arabidopsis*, gen “*Terminal Flowering 1 (TFL1)*”, “*LEAFY*” dan gen sasaran “*CONSTANS (CO)* ” meliputi gen “*FLOWERING*

*LOCUS T (FT)*” dan gen “*SUPPRESSOR OF OVEREXPRESSION CO 1 (SOC 1)*” memainkan peranan utama dalam mempromosi dan mengawal waktu pembungaan.

Dalam mengkaji perubahan identiti meristem dan pangawalan perubahan pembungaan di dalam kelapa sawit, kami memindahkan gen-gen *pCAMBIA/TFL1*, *pCAMBIA/JIT60*, *pCAMBIA/LFY*, *pGA/LFY* dan *pCAMBIA/SOC1* ke dalam sel embriogenik kelapa sawit. Penyelidikan ini memfokuskan kepada pengoptimuman transformasi menggunakan kaedah *Agrobacterium* dan penganalisaan tumbuhan transgenik. Objektif penyelidikan ini juga adalah untuk mengklon gen putatif *SOC1* dan *LFY* ke dalam vektor plasmid bersistem binari, mentransformasikan gen tersebut ke dalam sel embriogenik kelapa sawit dan seterusnya mengkaji ekspresi gen tersebut dengan menggunakan promoter CaMV35S.

Kejayaan pemindahan gen menggunakan *Agrobacterium* ke dalam genom tumbuhan sering bergantung kepada beberapa faktor termasuk suhu yang digunakan semasa ko-kultivasi, vektor binari dan promoter yang digunakan, dan juga jenis tumbuhan itu sendiri. Faktor yang paling penting yang boleh menghasilkan kejayaan untuk pemindahan T-DNA ke dalam genom tumbuhan adalah jenis bahan tumbuhan yang digunakan. Kami menggunakan sel embriogenik kelapa sawit terampai sebagai bahan untuk proses transformasi kerana kandungan sel ‘totipotent’ yang terdapat pada bahan tersebut.

Keputusan daripada penyelidikan menunjukkan, ampai sel embriogenik adalah sesuai untuk transformasi menggunakan *Agrobacterium*. Tumbuhan kelapa sawit transgenik yang stabil, di dalam tabung uji telah dihasilkan melalui kaedah pemindahan gen menggunakan *Agrobacterium* dalam tempoh 7 hingga 8 bulan selepas ko-kultivasi. Sejumlah 101 tumbuhan transgenik yang muncul daripada sekumpulan sel embriogenik diperolehi daripada 5563 kumpulan sel embriogenik. Kadar transformasi secara keseluruhan adalah 1.82%, tetapi ini adalah berdasarkan purata penghasilan daripada rawatan yang berbeza. Penganalisan menggunakan 'PCR' dan 'Southern Blotting' mengesahkan kemasukan T-DNA ke dalam genom tumbuhan kelapa sawit.



## ACKNOWLEDGEMENTS

I would like to express my most sincere gratitude and deepest appreciation to Assoc. Prof . Dr. K. Harikrishna for giving me the opportunity to persue my studies in the field of plant biotechnology. With his constant motivation and words of advice, has inspired me to become a better researcher and individual. Special thanks are extended to other members of my committee, Dr. Ho Chai Ling and Assoc. Prof. Dr. Ruslan Abdullah for their advice, comments and guidance whenever sought. To Ms. Girlie Wong (Applied Agriculture Research Sdn.Bhd), the deepest appreciation for the supply of tissue culture materials. My deepest appreciation also goes to my employer, the Felda Agriculture Service Sdn.Bhd, especially to Dr. Maheran Abu Bakar and Mr. Aw Khoo Teng for granting some time off, for me to complete my thesis.

Special thanks extended to the genetic lab members especially Pick Kuen, Choong, Yang Ping, Mei Chooi, See, Radziah, Weng Wah and etc. Thanks also to Mr.Ong and Kak Fadilah for all their help and support.

Finally, I wish to express my deepest gratitude and appreciation to my family for their constant support throughout my study. To my best friend, Tengku Adnin, thank you for your support and occasional constructive criticisms of my work.

I certify that an Examination Committee met on 27<sup>th</sup> June 2003 to conduct the final examination of Siti Habsah Roowi on her Master of Science thesis entitled “*Agrobacterium*-mediated transformation of Oil Palm (*Elaeis guineensis* Jacq.) Suspension Culture” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Norihan Mohd. Saleh, Ph.D.**

Associate Professor,  
Department of Biotechnology,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia.  
(Chairperson)

**Harikrishna Kulaveerasingam, Ph.D.**

Associate Professor,  
Department of Biotechnology,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia.  
(Member)

**Ruslan Abdullah, Ph.D.**

Associate Professor,  
School of Bioscience and Biotechnology,  
Faculty of Food Science and Technology,  
Universiti Kebangsaan Malaysia.  
(Member)

**Ho Chai Ling, Ph.D.**

Department of Biotechnology,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia.  
(Member)



**GULAM RUSUL RAHMAT ALI, Ph.D**

Professor/Deputy Dean  
School of Graduate Studies  
University Putra Malaysia

Date : 02 SEP 2003



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**Harikrishna Kulaveerasingam, Ph.D.**

Associate Professor,  
Department of Biotechnology,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia.  
(Chairman)

**Ruslan Abdullah, Ph.D.**

Associate Professor,  
School of Bioscience and Biotechnology,  
Faculty of Food Science and Technology,  
Universiti Kebangsaan Malaysia.  
(Member)

**Ho Chai Ling, Ph.D.**

Department of Biotechnology,  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia.  
(Member)



**AINI IDERIS, Ph.D.**  
Professor/Dean  
School of Graduate Studies  
University Putra Malaysia

Date : 15 SEP 2003

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledge. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

  
(SITI HABSAH ROOWT)

Date: 8.9.2003

## TABLE OF CONTENTS

	Page
ABSTRACT	II
ABSTRAK	V
ACKNOWLEDGEMENTS	VII
APPROVAL	IX
DECLARATION	X
LIST OF TABLES	XIV
LIST OF FIGURES	XV
LIST OF ABBREVIATIONS	XVIII

### CHAPTER

1	INTRODUCTION	1
2	LITERATURE REVIEW	4
2.1	The Botany of Oil Palm	4
2.2	Tissue Culture of Oil Palm	6
2.3	Abnormalities Derived from Tissue Culture	8
2.4	Somatic Embryogenesis	11
2.4.1	Oil Palm Embryogenic Suspension Culture	12
2.4.2	Transformation of Embryogenic callus	13
2.4.3	The Transition of Embryogenic Callus to Whole Plants	15
2.4.4	The Meristem Genes	18
2.5	<i>Agrobacterium tumefaciens</i>	22
2.5.1	The Transfer of T-DNA from <i>Agrobacterium tumefaciens</i> into plants	23
2.5.2	The Successful Transfer Of Genes Using <i>Agrobacterium tumefaciens</i>	26
2.5.3	Promoters	27
2.5.4	Selectable Markers	30
2.6	Oil Palm Transformation	33
3	MATERIALS AND METHODS	35
3.1	General methods for construction of plant transformation vectors	35
3.1.1	Preparation of Insert	35
3.1.2	Preparation of Vector	35
3.1.3	Large Scale Plasmid Preparation	35
3.1.4	Gel Electrophoresis of Plasmid DNA	37
3.1.5	Polymerase Chain Reactions	38



3.1.6	Purification of Restricted Plasmid	39
3.1.7	Purification of DNA from Agarose Gel	39
3.1.8	Ligation	40
3.1.9	Transformation	40
3.1.10	Screening of Transformants	40
3.1.11	Preparation of Competent cells of Disarmed <i>Agrobacterium</i> LBA 4404 and Transformation	41
3.1.12	<i>Agrobacterium</i> DNA extraction	42
3.2	Construction of Expression Vectors	42
3.2.1	Construction of Binary Vector for <i>SOC1</i> gene	47
3.2.2	Construction of Binary Vector for <i>LFY</i> gene	48
3.3	Transformation of Embryogenic callus	49
3.3.1	Tissue Culture Materials	49
3.3.2	Tissue Culture Media and Chemicals	49
3.3.3	<i>Agrobacterium</i> with Binary Vectors	49
3.3.4	<i>Agrobacterium</i> Culture Medium and Antibiotics	51
3.3.5	Determination of Minimal Inhibitory Concentration of Hygromycin	51
3.3.6	Growth of <i>Agrobacterium</i>	51
3.3.7	Embryogenic Callus Treatment	52
3.3.8	Plant Regeneration	54
3.3.9	Analysis of Transgenic Oil Palm	54
4	RESULTS	59
4.1	Construction of Plant Transformation Vectors	59
4.1.1	Construction of Binary Vector for <i>SOC1</i> gene and transformation of Binary vector into <i>Agrobacterium</i> <i>tumefaciens</i> LBA 4404	59
4.1.2	Construction of Binary Vector for <i>LFY</i> gene and Transformation of Binary Vector into <i>Agrobacterium tumefaciens</i> strain LBA 4404	65
4.2	Plant Transformation	70
4.2.1	Minimal Inhibitory Concentration of Hygromycin	70
4.2.2	The <i>Agrobacterium tumefaciens</i> Growth Curve	72
4.2.3	Inoculation and Co-Cultivation of Embryogenic Callus with <i>Agrobacterium</i> and production of transgenic plant	73
4.2.4	Histochemical analysis of GUS expression	76
4.2.5	The Effect of Temperature during Co-Cultivation	77

4.2.6	The Effect of Different Binary Vectors used for Transformation	79
4.2.7	The Effect of Genotype in Oil Palm Transformation	81
4.2.8	PCR and Southern Blot Analysis	82
5	DISCUSSION	88
6	CONCLUSION	94
	BIBLIOGRAPHY	96
	APPENDICES	106
	VITAE	115

## LIST OF TABLES

Table		Page
1	Example of different type of plant promoters used for construction of transgene	29
2	Antibiotics selection markers used for plant transformation	31
3	Herbicide selection markers used for plant transformation	32
4	Metabolic selection markers used for plant transformation	32
5	Commonly used reporter genes in plant transformation	33



## LIST OF FIGURES

Figure		Page
1	The process of seed germination	5
2	The oil palm tissue culture process	7
3	Root apical meristem	15
4	Shoot apical meristem structure	16
5	Schematic of a section through the vegetative SAM of a dicotyledonous Plant	17
6	The signaling pathways that promote flowering in <i>Arabidopsis</i>	20
7	Basic step in the transformation of plant cells by <i>Agrobacterium tumefaciens</i>	26
8	<i>AoPR1-GUS</i> expression in tobacco	28
9	Plasmid map for pHK17	44
10	Illustration of Plasmid map of pHK17 digested with <i>Pst</i> I and <i>Bgl</i> II	44
11	The schematic layout for the construction of transformation vector pCAMBIA/SOC1	45
12	The schematic layout for the construction of transformation vector pCAMBIA/LFY	46
13	A schematic diagram demonstrating the construction of binary vector pCAMBIA/SOC1	47
14	A schematic diagram demonstrating the construction of binary vector pCAMBIA/LFY	48

15	The schematic diagram for the transformation of embryogenic suspension cultures	50
16	Double digestion of pSH2 with <i>EcoRI</i> and <i>XbaI</i> and electrophoresed on 1% (w/v) agarose gel	60
17	Double digestion of pSH3 with <i>EcoRI</i> and <i>XbaI</i> and electrophoresed on a 1% (w/v) agarose gel	61
18	Orienting inserts in pSH1 construction	62
19	Analysis of recombinant pCAMBIA1300/SOC1 with <i>EcoRI</i> and <i>XbaI</i>	63
20	Confirmation of pCAMBIA1300/SOC1 into <i>Agrobacterium</i> LBA 4404	64
21	Plasmid map of pCAMBIA/SOC1	64
22	Gel electrophoresis of inserts after purification process	65
23	Confirmation of integrating of pSH4 construction	66
24	Orienting inserts in pSH4 construction	66
25	Confirmation of <i>pCAMBIA1300/LFY</i> with <i>EcoRI</i> and <i>XbaI</i>	67
26	Transformation of pCAMBIA1300/LFY into <i>Agrobacterium</i> LBA 4404 and confirmed by PCR	68
27	PCR confirmation of pGA/LFY	68
28	Plasmid map of pCAMBIA/LFY	69
29	Plasmid map of pGA/LFY	69
30	Determination of optimal concentration of hygromycin for selection	71
31	<i>Agrobacterium tumefaciens</i> (LBA 4404) growth curves	73
32	Studies in the production of transgenic Plants	75
33	Histochemical analysis of GUS expression	76

34	The formation of embryoid after 5 weeks on MS basal medium + 500 mg/l carbenicillin	78
35	Frequency of embryoid formation at different incubation temperature during co-cultivation	78
36	Induction of embryoid on selection medium by using two different binary vectors	79
37	Transformation frequency using 2 types of binary vectors pCAMBIA/LFY and pGA/LFY	80
38	Plant regeneration frequency from four different genotype of embryogenic suspension of oil palm	81
39	Analysis of putative transgenic plants carrying <i>Terminal flowering gene 1 (tfl1)</i>	83
40	Analysis of putative transgenic plants carrying the <i>LEAFY</i> gene ( <i>lfy</i> )	84
41	Analysis of putative transgenic plants carrying the <i>SOC1</i> ( <i>Suppressor of Overexpression 1</i> ) gene	85
42	Analysis of putative transgenic plants carrying the reporter gene ( <i>GUS</i> )	86
43	Confirmation of transgene integration of transgenic oil palm plants carrying the <i>terminal flowering 1</i> gene	87

## LIST OF ABBREVIATIONS

ANOVA	analysis of variance
AS	acetosyringone
bp	base pair
CaMV	cauliflower mosaic virus
CIM	callus induction medium
°C	degree centigrade
DNA	deoxyribonucleic acid
2,4-D	2,4-dichlorophenoxyacetic acid
DMSO	dimethyl sulfoxide
DMRT	Duncan's Multiple Range Test
EDTA	ethylenediaminetetraacetic acid
GUS	$\beta$ -Glucuronidase
g	gram
HPT	hygromycin phosphotransferase gene
h	hour (s)
mM	millimolar
$\mu$ M	micromolar
ml	milliliter
mg	milligram (s)
MS	Murashige and Skoog
MSO	hormoneless Murashige and Skoog
NaOH	Sodium hydroxide

Nos ter	Nopaline synthase gene terminator
NPTII	Neomycin phosphotransferase gene
OD	Optical density
%	Percentage
PCR	Polymerase Chain Reaction
RNase	ribonuclease
rpm	revolution per minute
35S	35 subunit of ribosome
T-DNA	transferable-DNA
$\mu$ l	microliter
v/v	volume per volume
w/v	weight per volume
X-gluc	5-bromo-4-chloro-3-indoyl-glucuronide

## **CHAPTER 1**

### **INTRODUCTION**

Oil palm is one of the larger members of the palm species. It has a single stem, and unlike some palms, does not produce suckers from the base. There are about 212 genera and about 2500 species of palms that grow in the tropics and subtropics. The single vegetative growing point is situated in a depression at the stem apex. The meristem is continuously active, producing a new leaf primordium about every two weeks in a mature palm. The leaf is pinnate, with the pinnae (leaflets) arranged in two or more planes on each side of the rachis. In each leaf axil produced in cycles of variable duration, are inflorescences that following pollination develop into a fruit bunch, from which mesocarp and kernel oils can be extracted.

The oil palm is a particularly important species in world trade as a major economic oil crop. It also has wide usage in the food and non-food industries. Currently, Malaysia is the largest palm oil producer in the world with 3.5 million hectares of land devoted to oil palm cultivation, representing more than 60% of the country's total cultivated agricultural land and has become the world's largest exporter of palm oil with a market share of 60.6%. However, soon Indonesia will overtake Malaysia and become the largest palm oil producer in the world. Thus, the palm oil industry in Malaysia is undergoing a challenging time.

Due to the increased demand for palm oil, increasing the oil yield is the main agenda for those involved in the oil palm industry. Research in plant genetics and breeding

has largely improved the performance of the crop, however, with conventional breeding and seed production methods the maximum potential of the selected hybrid genotypes may still not be realized. Therefore, vegetative propagation of elite hybrid oil palm via tissue culture seems to be an attractive alternative.

Gene transfer to elite germplasm is an attractive method for more rapid genetic improvement of this perennial crop. Several techniques such as direct DNA uptake using biolistic or particle gun bombardment, and vector-mediated gene delivery using *Agrobacterium tumefaciens*, have been used for oil palm transformation. However, the success of gene transfer depends on several factors, such as the use of suitable promoters and the appropriate selectable marker gene with a sensitive selection agent, the use of appropriate target tissue and development of a reproducible regeneration protocol. The lack of an efficient regeneration method is a great bottleneck in the application of transformation technology to oil palm. The regeneration of genetically transformed plants based on somatic embryogenesis could be a viable solution for the oil palm.

The aim of this thesis was to establish an efficient procedure for the stable transformation of foreign genes into oil palm via co-cultivation with *A. tumefaciens* and to study the integration and expression of foreign genes in transgenic oil palm by molecular analyses such as PCR and Southern blot hybridization. The experiments conducted in this thesis are divided into 3 parts:

1. Construction of binary transformation vector.
2. Transformation of several genes of interest into oil palm embryogenic suspension cultures.
3. Analysis of putative transformed plantlets.



## CHAPTER 2

### LITERATURE REVIEW

#### 2.1 The Botany of Oil Palm

The oil palm (*Elaeis guineensis* Jacq.) is classified under the family Palmae in the order Palmales and grouped with the cocos, coraza and other genera under the tribe Cocoinae. It is a large feather palm having a solitary columnar stem with short internodes. The palm is normally monoecious with male or female, or at certain stages hermaphrodite inflorescences developed in the axil of the leaves. The fruit is a drupe, which is borne on a large compact bunch. The oil palm seed is a nut that remains after the soft oily mesocarp has been removed. It is composed of a shell or endocarp surrounding usually one kernel and sometimes up to 4 kernels. The embryo is straight and about 3mm in length. Its distal end lies opposite the germ pore separated by a thin layer of endosperm cells.

The seedling has three months to establish itself as an organism capable of photosynthesis and absorption of nutrients from the soil. The first adventitious roots are produced in a ring just above the radicle hypocotyl junction and these give rise to secondary roots before the first foliage leaf has emerged. The radicle continues to grow for about six months by which time it would have reached about 15 cm in length (Figure 1).